

Pythagorean Triangle With (One leg of right triangle $p^2 - q^2$) $- 2 \frac{\text{area}}{\text{perimeter}} = \alpha^3 - \beta^3$

S. Sriram¹, P. Veeramallan²

¹Assistant Professor, P. G & Research Department of Mathematics, National College, Tiruchirappalli, Tamilnadu, India

²Post Graduate Assistant in Mathematics, GHSS, Perumangalam, Villupuram, Tamilnadu, India

ABSTRACT

Pattern of Pythagorean Triangle for each of which, the ratio (One leg of right triangle ($p^2 - q^2$)) $- 2 \left(\frac{\text{Area}}{\text{Perimeter}} \right)$ may be expressed as the difference to two cubes as integers. A few interesting relations are also given.

Keywords : Pythagorean triangle, Ratio (Area/perimeter) as Difference of Two Cubic Integers.

I. INTRODUCTION

It is well known that Pythagoras triangle is a treasure house which contains many interesting results for an extensive review of the literature. The method of obtaining three non-zero integers x, y and z under certain relations satisfying the equation $x^2 + y^2 = z^2$ has been a matter of interest to various mathematicians. One may refer [1-7]. In [8-11] special Pythagoras problems are studies. In this communication we, present yet another interest Pythagorean Triangles, where in each of which, (One leg of right triangle ($p^2 - q^2$)) $- 2 \left(\frac{\text{Area}}{\text{Perimeter}} \right)$ may be expressed as the difference to two cubes as integers. A few interesting relations are also given. In addition, the recurrence relations for the sides of the triangle are presented.

Notation

$$t_{m,n} = n \left[1 + \frac{(n-1)(m-2)}{2} \right] = \text{poligonal number of rank } n \text{ with sides } m$$

II. METHODS AND MATERIAL

Method of Analysis

The most cited solution of the Pythagorean equation $x^2 + y^2 = z^2$ is

$$\begin{aligned} x &= 2pq \\ y &= p^2 - q^2 \text{ where } p > q > 0 \\ z &= p^2 + q^2 \end{aligned} \quad (1)$$

Denoting the Area and the Perimeter of the above Pythagorean triangle by A and P respectively, the assumption that the ratio $\frac{A}{P}$ can be expressed as the difference of two cubic non zero integers leads to the equation

$$p(p - q) = \alpha^3 - \beta^3 \quad (2)$$

Where α, β are the non zero integer. ($\alpha > \beta > 0$)
Choosing

$$p = \alpha^2 + \beta^2 + \alpha\beta, \quad p - q = \alpha - \beta \quad (3)$$

in (2), and solving we get, $p = \alpha^2 + \beta^2 + \alpha\beta$ and

$$q = \alpha^2 + \beta^2 + \alpha\beta - \alpha + \beta \quad (4)$$

In which follows, we obtain the values of x , y , z

III. REFERENCES

$$x(\alpha, \beta) = 2pq \implies x = 2\alpha^4 + 2\beta^4 + 6\alpha^2\beta^2 + 4\alpha\beta^3 + 4\alpha^3\beta + 2\beta^3 - 2\alpha^3 \quad (5)$$

$$y(\alpha, \beta) = p^2 - q^2 \implies y = 2\alpha^3 - 2\beta^3 + 2\alpha\beta - \alpha^2 - \beta^2 \quad (6)$$

$$z(\alpha, \beta) = p^2 + q^2 \implies z = 2\alpha^4 + 2\beta^4 + 6\alpha^2\beta^2 + 4\alpha\beta^3 + 4\alpha^3\beta + 2\beta^3 - 2\alpha^3 + 2\beta^3 - 2\alpha\beta + \alpha^2 + \beta^2, \text{ where } \alpha > \beta \quad (7)$$

Few examples are given

α	β	p	q	x	y	z
2	1	7	6	88	13	85
4	3	37	36	2664	73	2665
5	4	61	60	7320	121	7321
9	7	193	191	73726	768	73730
3	2	19	18	684	37	685
7	3	79	75	11850	616	11816
5	2	39	36	2808	225	2817
7	4	93	90	16740	549	16749
8	5	129	126	32508	765	32517
7	2	67	62	8308	645	8333

Properties

- $x(\alpha, \beta) \equiv 0 \pmod{2}$
- $x(\alpha, 1) - 4\alpha Pen_\alpha - Obl_\alpha + 2\alpha$ is a perfect square.
- $6(x(\alpha, 1) - 4\alpha Pen_\alpha - Obl_\alpha + 2\alpha)$ is a nasty number.
- $(x - z) + (\alpha^2 + \beta^2) \equiv 0 \pmod{2}$
- $6(z - x)$ is a nasty number.
- $y(\alpha, \beta) + (\alpha^2 + \beta^2) \equiv 0 \pmod{2}$
- $z(\alpha, 1) = (T_{4,\alpha} + 1)(4T_{3,\alpha} + 5)$
- $z(\alpha, 1) = 4(T_{4,\alpha})(T_{3,\alpha}) + 2T_{9,\alpha} + 7\alpha + 5$
- $y(\alpha, 1) - (\alpha^2 + 1) \equiv 0 \pmod{2}$
- $z(\alpha, \beta) - (\alpha^2 + \beta^2) \equiv 0 \pmod{2}$
- $z(\alpha, 1) - 4\alpha Pen_\alpha - 12Tri_\alpha - 5$ is a perfect square.
- $x(\alpha, \beta) - (y(\alpha, \beta) + z(\alpha, \beta)) \equiv 0 \pmod{2}$
- $x(\alpha, \beta) - (y(\alpha, \beta) + z(\alpha, \beta)) + 2(\alpha^3 - \beta^3)$ is a perfect square
- $y(\alpha, \beta) + (\alpha^2 + \beta^2) \equiv 0 \pmod{2}$
- $y + z = 0 \pmod{2}$
- $x - (y + z) = 0 \pmod{2}$

- [1] Dickson, L.E., History of the theory of numbers, Vol. II, Chelsea Publishing Company, New York (1952).
- [2] Smith, D.E., History of Mathematics, Vol. I and II, Dover Publications, New York (1953).
- [3] Boyer, Carl, B. and Merzbach, U.T.A.C., A History of Mathematics, John Wiley and Sons (1989).
- [4] Akituro, Nishi, A method of obtaining Pythagorean Triples, Amer. Math. Monthly, Vol. 94, No. 9, 869-871 (1987).
- [5] Albert H. Beiler, "Recreations in the Theory of Numbers", Dover Publications, New York, 1963.
- [6] S. B. Malik, "Basic Number Theory", Vikas Publishing House Pvt. Limited, New Delhi. 1998.
- [7] L. J. Mordell, "Diophantine Equations", Academic Press, New York, 1969.
- [8] S.G. Telang, "Number Theory", Tata McGraw-Hill Publishing Company, New Delhi, 1996.
- [9] Thomas Koshy, "Elementary number Theory with Applications", Academic Press, 2005.
- [10] T.Nagell, "Introduction to Number Theory", plencem, New York, 1969.
- [11] Gopalan, M.A., Note on Integral Solutions of $X^2 + Y^2 = Z^2$, ActaCienciaIndica, Vol. XXVII M, No. 4, 493 (2001)
- [12] M. A. Gopalan and S. Leelavathi "Pythagorean Triangle with $2(\text{Area} / \text{Perimeter})$ as a Cubic Integer", Bulletin of Pure and Applied Sciences, Vol. 27, No. E(2),2007, pp. 197-200.
- [13] M. A. Gopalan and G. Janaki "Pythagorean Triangle with Area / Perimeter as a special polygonal number", Bulletin of Pure and Applied Sciences, Vol.27, No.E(2),2007, pp.393-402.
- [14] M. A. Gopalan and S. Devibala, Pythagorean Triangle: A Treasure House, Proceeding of the KMA National Seminar on Algebra, Number Theory and Applications to Coding and Cryptanalysis, Little Flower College, Guruvayur, September 16-18, 2004
- [15] M. A. Gopalan and S. Devibala, "Pythagorean Triangle with Triangular number as a leg", Impact J.Sci.Tech.vol.2(4),2008,pp.195-199..
- [16] M. A. Gopalan and S. G.Janaki, "Pythagorean Triangle with Nasty number as a leg", Journal of Applied Mathematical Analysis and Applications", vol.4(1-2),2008,pp.13-17.

- [17] M. A. Gopalan and S. G. Janaki, "Pythagorean Triangle with Perimeter as a pentagonal number", *Antarctica J. Math.*, vol.5(2), 2008, pp.15-18.
- [18] M. A. Gopalan and G. Gnanam "Pythagorean Triangles and special polygonal numbers", *International Journal of Mathematical Sciences* vol.9(1-2), 2010, pp.211-215.
- [19] M. A. Gopalan and G. G. Sangeetha, "Pythagorean Triangles with Perimeter as Triangular number", *Global Journal of Applied Mathematics and Mathematical Sciences*, vol.3(1-2), 2010, pp.93-97.
- [20] M. A. Gopalan and B. Sivakami, "Pythagorean Triangle with Hypotenuse minus 2(Area / Perimeter) as a Square Integer", *Archimedes J. Math.*, vol.2(2), 2012, pp. 153-166.
- [21] M. A. Gopalan and V. Geetha, "Pythagorean Triangle with Area / Perimeter as a special polygonal number", *International Refereed Journal of Engineering and Science*, vol.2(7), 2013, pp.28-34.
- [22] M. A. Gopalan and Manju Somanath and K. Geetha, "Pythagorean Triangle with Area / Perimeter as a special polygonal number", *IOSR-JM*, vol.7(3), 2013, pp. 52-62.
- [23] M. A. Gopalan and Manju Somanath and V. Sangeetha, "Pythagorean Triangles and pentagonal number", *Cayley J. Math*, vol.2(2), 2013, pp.151-156.
- [24] M. A. Gopalan and Manju Somanath and V. Sangeetha, "Pythagorean Triangles and special pyramidal numbers", *IOSR-JM*, vol.7(4), 2013, pp. 21-22.
- [25] P. Thirunavukkarasu and S. Sriram, "Pythagorean Triangle with Area / Perimeter as quartic integer", *International Journal of Engineering and Innovative Technology (IJEIT)*, vol.3(7), 2014, pp.100-102.
- [26] M. A. Gopalan and R. Anbuselvi, "A Special Pythagorean Triangle", *Acta Ciencia Indica XXXI M*, No. 1, p. 053, 2005.
- [27] M. A. Gopalan and S. Devibala, "On a Pythagorean Problem", *Acta Ciencia Indica, XXXII M*, No. 4, p. 1451, 2006.
- [28] S. Sriram, "Pythagorean Triangle with (one Leg of Right Triangle $p^2 - q^2$) - $2(\text{area/perimeter}) = \alpha\beta$ ", *International Journal of Engineering and Innovative Technology, IJEIT*, vol. 6, Issue. 2, August 2016.
- [29] M. A. Gopalan and S. Leelavathi "Pythagorean Triangle with Area / Perimeter as a Square Integer", *International Journal of Mathematics, Computer Sciences and Information Technology* Vol. 1, No. 2, July-December 2008, pp. 199-204.
- [30] M. A. Gopalan and J. Kaliga Rani, "A Special Pythagorean Triangle", *Acta Ciencia Indica, XXXII M*, No. 4, p. 1451, 2006.