

Use of 2-Hydroxy-4-Isobutoxy Acetophenone Oxime as Spectrophotometric Reagent for Copper

Dr. J. R. Shukla

Associate Professor (Principal), Chemistry Department, Shri C.N.P.F Arts & D.N.Science College, Dabhoi, Gujarat, India

ABSTRACT

2-Hydroxy-4-isobutoxy acetophenone oxime(HIBAO) has been used for the spectrophotometric determination for Cu(II) at pH range 7 to 9 in chloroform medium. Job's method for continuous variation, Yoe and Jones' mole ratio method, the slope ratio method show metal ligand ratio in complex to be 1:2. The stability constant of the complex is found to be 4.32×10^8 . The dark green coloured complex obeys Beer's law in the concentration range 31.77 to 444.78 ppm for Cu(II) ion, while the optimum concentration range from Ringbom plot is found to be 127.08 to 381.24ppm. The photometric sensitivity and molar absorptivity at the 650nm are found to be $0.488 \mu\text{g}/\text{cm}^2$ and $130 \text{ l mol}^{-1}\text{cm}^{-1}$ respectively. The standard free energy of formation of complex is -12.00 kcal/mole at 30°C . The complex is stable for 48 hrs. The reagent has also been found to give quite satisfactory results for Cu(II) present in alloy like brass, bronze and synthetic mixtures. The antimicrobial Activity of HIBAO and Cu-HIBAO complex have also been checked.

Keywords: 2-Hydroxy-4-Isobutoxy Acetophenone Oxime, IBAO, Systronic pH meter

I. INTRODUCTION

Various o-hydroxy phenones, phenone oximes, phenyl hydrozones, chalknoneoximes, etc, have been used as an analytical reagent for the spectrophotometric and gravimetric determination of copper and other transition metal ions¹⁻⁵. In the present work the use of 2-hydroxy-4-isobutoxy acetophenone oxime (HIBAO) as photometric reagent for Cu(II) has been described.

II. METHODS AND MATERIAL

A 0.1M stock solution of Cu(II) has been prepared by dissolving copper sulphate (A.R) in distilled water containing few drops of sulphuric acid. The amount of Cu(II) in this solution was determined following standard procedures⁶.

Preparation of 2-hydroxy-4-isobutoxy acetophenone oxime (HIBAO):

Resacetophenone was prepared from resorcinol by standard methods⁷. 2-hydroxy-4-isobutoxyacetophenone(HIBA) has been prepared by refluxing resacetophenone and isobutyl bromide in suitable solvent for 4 hrs. 2-hydroxy-4-isobutoxyacetophenoneoxime(HIBAO)⁸ has been prepared by refluxing HIBA with hydroxyl amine hydrochloride in the presence of sodium acetate in ethanol medium for 4 hrs. The reagent when recrystallized from ethanol was obtained in the colourless, needle like crystals (m.p. $102^\circ\text{C} \pm 1^\circ\text{C}$), with M.W 223.40 (calcd. for $\text{C}_{12}\text{H}_{17}\text{NO}_3 = 223.27$). The reagent is insoluble in water but soluble in alcohol, acetone benzene chloroform, carbon tetrachloride, etc. The elemental analysis and spectral analysis of the compound confirm its structure.

Preparation of Cu(II)-HIBAO complex and selection of solvent : When an alcoholic solution of HIBAO was added to 0.01 M aqueous metal ion

solution, dark green precipitates of complex were obtained in the pH range 3.0 - 10.0. The complex was found to be insoluble in polar solvents like water, methanol or ethanol but soluble in non-polar solvents like chloroform, benzene, CCl₄ etc. As Cu-HIBAO complex was more soluble in chloroform, it was selected as a solvent for extractive spectrophotometric determination of Cu(II).

Apparatus : Spectrophotometric measurements were made with a Systronics UV/VIS spectrophotometer (model-118) using 10mm glass cells. All pH measurements were made with Systronic pH meter (model-324).

III. RESULTS AND DISCUSSION

Results are given in Fig. I-IV and Table I-II.

Table I: Stability Constant of Cu(II) –HIBAO Complex at 30°C

Method employed	Em	Es	α	K (n=1)
Mole ratio method				
Set-I	0.337	0.322	0.04451	4.33×10^8
Set-II	0.179	0.166	0.07263	3.87×10^8
Job's Method				
	0.260	0.247	0.05000	4.75×10^8
Set-I	0.520	0.503	0.03269	4.33×10^8
Set-II				
Mean K Stab				4.32×10^8

Table II: Analysis of copper in various samples

Sample	"Cu" taken		Absorbance	"Cu" found		Relative error(%)
	μg	%		μg	%	
Brass	988.8	61.80	0.201	956.7	60.58	1.97
			0.206	980.1		
			0.208	970.9		
			Avg.	969.3		
Bronze	1411.2	88.20	0.288	1370.8	87.56	0.73
			0.296	1408.8		
			0.299	1423.1		
			Avg.	1400.9		
Synthetic mixture No.1	2541.6	-	0.511	2497.6	-	0.51
			0.522	2551.4		
			0.535	2614.9		
			Avg.	2554.6		
Synthetic mixture No.2	953.1	-	0.198	967.8	-	0.84
			0.191	933.5		
			0.201	982.4		
			Avg.	961.2		
Synthetic mixture No.3	2223.9	-	0.457	2233.7	-	0.88
			0.428	2091.9		
			0.468	2287.4		
			Avg.	2204.3		

Optimum pH and Selection of Wavelength : The pH of the solution has pronounced effect on the reaction between Cu(II) and HIBAO and the stability of the complex. On the other hand the absorbance is dependant upon the wavelength used. Both the parameters were therefore controlled to give maximum absorbance. Absorbance measurements of the reagent in chloroform show maxima at 240nm, 268nm and 301nm with negligible absorbance beyond 400nm. The absorbance measurements of Cu(II)-HIBAO complex show a maxima at 400nm and 650nm. As the interference due to the reagent appeared to be negligible a wavelength of 650 nm was selected for the present work.

To determine the optimum pH for complex formation series of buffer solutions with pH values ranging from 3.0 to 10.0 were prepared. To above buffer solutions, 2.0 ml of 0.01 M Cu(II) solution and 10 ml 0.01 HIBAO solution in chloroform were added. After shaking the mixture for two minutes, the dark green coloured complex was extracted. The absorbance of organic layer containing complex was measured at 650nm against a blank. From the results given in Fig. I, it may be generalized that maximum absorbance takes place at pH range 7.0 to 9.0. Hence a pH of 8.0 and wavelength of 650nm have been selected for the present work.


Reproducibility : Absorbance measurements of a set of ten solution prepared in a similar way and containing the same concentrations of all the reagents show that the reproducibility of measurements are quite good with a standard deviation of ± 6.289 Units, i.e., 0.25%.

Effect of time and temperature : To determine the effect of time and temperature on the intensity of colour and the stability of the Cu(II)- HIBAO complex, absorbance was measured at room temperature (30°C) at regular intervals of time up to 48 hrs and also at temperatures of 30°C to 55°C. The results show that complex is stable ($\pm 2\%$ deviation) for two weeks and up to 45°C.

Stoichiometry and Stability Constant of the Complex : The method of Vosbough and Cooper⁹ showed that one complex is formed. To determine the stoichiometry of complex, Yoe and Jones mole ratio method¹⁰ the slope ratio method¹¹ and Job's method of continuous variation¹² were employed (Fig. II-IV). All the three methods show a 1:2 metal:ligand ratio in the complex.

The value of the stability constants calculated from the job's method as well as from the mole ratio method are given in Table-I. From the table the average value of stability constant may be taken as 4.32×10^8 . The standard free energy of formation of the complex, ΔG° , is -12.00 kcal/mole at 30°C.

The IR spectras of reagent and complex revealed that the -OH (stretch) band of 3393cm^{-1} for the reagent disappears when the complex is formed i.e., the complex formation takes place through the N of oximino group and O- of the 2-hydroxy group. Based on above data the Cu(II)-HIBAO complex can be assigned the following structure.



Conformity to Beer's law and the optimum concentration range : Beer's law is obeyed between the range 31.77 to 444.78ppm of Cu(II). At higher concentrations negative deviations occur. The optimum concentration range for determination of Cu(II) in solution, as deduced from the Ringbom plot¹³, is found to be 127.08 to 381.24ppm. The molar absorptivity (ϵ) of the complex is $130 \text{ mol}^{-1}\text{cm}^{-1}$ and the photometric sensitivity as per Sendell's definition¹⁴ is found to be $0.488 \mu\text{g}/\text{cm}^2$ at 650nm.




Figure I : Effect of pH




Figure II : Job's method of continuous variation
 Set-I : 0.01 M Cu(II) and 0.01 M HIBAO
 Set-II : 0.02 M Cu(II) and 0.02 M HIBAO




Figure III : Yoe and Jones mole ratio method
 Set-I : 0.005M Cu(II) and 0.010M HIBAO,
 Set-II : 0.0025M Cu(II) and 0.005M HIBAO




Figure IV : Slope ratio method.
 10 ml 0.01 M HIBAO ; 0.01M Cu(II) (variable) 10
 ml 0.01 M Cu(II) ; 0.01 M HIBAO (variable)

Effect of diverse ions : The interference due to the presence of other ions on the determination copper ions as Cu(II)-HIBAO complex has also been studied. A difference of more than $\pm 2\%$ in the absorbance value has been considered as interference. According to this criterion, the tolerance limits of various ions, expressed in μg , for a solution containing $1270.80 \mu\text{g}$ (Cu(II)) are as follows.

- up to 100000 μg : $\text{Na}^+, \text{K}^+, \text{NH}_4^+, \text{Cl}^-, \text{NO}_3^-$,
 $\text{SO}_4^{2-}, \text{CH}_3\text{COO}^-$
- up to 10000 μg : $\text{Ca}^{+2}, \text{Ba}^{+2}, \text{Sr}^{+2}, \text{Mg}^{+2}, \text{Al}^{+3}$,
 $\text{Zn}^{+2}, \text{Cd}^{+2}$,
 MoO_4^- , citrate, tartrate,
oxalate
- up to 1000 μg : $\text{Cr}^{+3}, \text{UO}_2^{+2}, \text{Pd}^{+2}$
- up to 100 μg : $\text{Ni}^{+2}, \text{Co}^{+2}, \text{Mn}^{+2}, \text{Fe}^{+3}$
- up to 10 μg : $\text{EDTA}, \text{S}_2\text{O}_3^{-2}$

Determination of copper from various samples : To determine the usefulness of the reagent in estimation of copper from various samples containing copper were taken and estimated by HIBAO. For this purpose, the alloy samples containing copper metal were dissolved in 1:1 nitric acid by heating on a sand bath. The resulting solution was made up to 250 ml with distilled water in a volumetric flask. The synthetic mixtures containing copper metal were also taken for analysis. Aliquot of this sample solution was pipeted out and its spectrophotometric determination was carried out by the proposed method. The result are given in Table-II.

Antimicrobial Activity :

HIBAO and Cu-HIBAO were screened for their antibacterial activity against Gram positive bacteria i.e. *Bacillus subtilis* and Gram negative bacteria i.e. *Pseudomonas aeruginosa* using Cup-plate agar diffusion method. The Fungi *Aspergillus niger* was used for antifungal activity of above compound using Cup-plate agar diffusion method. HIBAO exhibit good antibacterial activity and excellent antifungal activity with respect to standard drugs while the Cu-HIBAO complex found quit good antibacterial and antifungal activity.

IV. ACKNOWLEDGEMENTS

The authors are thankful to Uttar-Purva Gujarat Uchcha Kelvani Mandal, Pilvai, for laboratory facilities. Thanks are also due to SICART, V.V Nagar, for providing C, H and N analysis.

V. REFERENCES

- [1]. Talati, J.D and K.S.Parikh, "Anarta", J. North Gujarat Uni., 3, 93, (1992).
- [2]. Talati, J.D and K.S.Parikh, J. Inst. Chemists (INDIA), 65, 84, (1993).
- [3]. Talati, J.D and K.S.Parikh, Asin J. Chem., 6, (1994).
- [4]. J.A.dave and S.S.Shah,Asian J. Chem., 20, 4141,(2008).
- [5]. Suketu I. Desai, Vikas A. Desai and H.B.Niak, Asian J. Chem.,20,4919,(2008).
- [6]. A.I.Vogel, "A Text Book of Quantitative Inorganic Analysis", 4th Ed., (Longaman, London), 1978, p.379.
- [7]. Brewster, C.M. and J.C.Harris, J.Amer Chem Soc., 52, 4866 (1930).
- [8]. A.I.Vogel, "A Text Book of Practical Organic Chemistry, 3rd Ed., (Longaman, London), p. 341 (1966).
- [9]. W.C.Vosburgh and G.R.Cooper, J.Amer.Chem.Soc., 63, 473 (1941).
- [10]. J.H.Yoe and A.L.Jones, Ind.Eng.Chem., Anal. Ed., 16, 111 (1944).
- [11]. Harvey, A.E. and D.L.Manning, J.Amer.Chem.Soc., 72, 4488 (1950).
- [12]. P.Job, Ann. Chem., 9, 113 (1928).
- [13]. Ringbom, Anal. Chem., 115, 332 (1938).
- [14]. E.B.Sandell, "Colorimetric determination of traces of metals", 3rd Ed., (inderscience, New York), pp.83, (1959).